



# Exclusive breast-feeding and sociodemographic characteristics are associated with dietary patterns in children aged 4–7 years

Sarah Aparecida Vieira<sup>1,\*</sup>, Poliana Cristina de Almeida Fonseca<sup>1</sup>, Cristiana Santos Andreoli<sup>2</sup>, Helen Hermana Miranda Hermsdorff<sup>1</sup>, Andréia Queiroz Ribeiro<sup>1</sup>, Patrícia Feliciano Pereira<sup>1</sup>, Silvia Eloiza Priore<sup>1</sup> and Sylvia do Carmo Castro Franceschini<sup>1</sup>

<sup>1</sup>Department of Nutrition, Federal University of Viçosa, Avenue Ph Rolfs, University Campus, Viçosa – MG, 36570-900, Brazil; <sup>2</sup>Department of Technology in Gastronomy, Federal Institute of Minas Gerais, Ouro Preto, MG, Brazil

Submitted 8 May 2017; Final revision received 5 October 2018; Accepted 22 October 2018

## Abstract

**Objective:** To identify the dietary patterns of children aged 4–7 years and verify their association with sociodemographic characteristics, lifestyle habits and exclusive breast-feeding (EBF).

**Design:** A cross-sectional study nested within a cohort, performed with Brazilian children aged 4–7 years. The children were re-evaluated at age 4 to 7 years and food patterns were identified *a posteriori* through principal component analysis. The predictive variables were related to socio-economic characteristics, lifestyle habits and duration of EBF.

**Setting:** Viçosa, Minas Gerais, Brazil.

**Participants:** Representative sample of 403 children followed up by the Lactation Support Program from the Extension Program of the Universidade Federal de Viçosa during the first 6 months of life.

**Results:** Five dietary patterns were identified: ‘Traditional’, ‘Unhealthy’, ‘Milk and chocolate’, ‘Snack’ and ‘Healthy’. Children who did not receive EBF until they were at least 4 months old had a higher adherence to the ‘Unhealthy’ and ‘Snack’ patterns, and older children also consumed more ‘Unhealthy’ foods. The highest income was associated with the highest consumption of foods of the patterns ‘Unhealthy’, ‘Milk and chocolate’ and ‘Healthy’.

**Conclusions:** In view of the results, we emphasize the importance of providing support and encouragement towards EBF in the first months of life, as it can positively influence lifelong eating habits.

**Keywords**  
Dietary patterns  
Exclusive breast-feeding  
Socio-economic status  
Children  
Cohort study  
Brazil

Childhood is the time of life at which there is the greatest tendency to develop inappropriate eating habits. Children are influenced by the family’s diet and lifestyle and are subject to changes in environmental and behavioural patterns due to the transition into the school environment<sup>(1)</sup>. There is evidence that dietary habits developed during this stage tend to be maintained<sup>(2,3)</sup> and influence lifelong health<sup>(4,5)</sup>.

Because of changes in population dietary habits, especially since the 1990s, with high consumption of ultra-processed products, the intake of foods rich in simple carbohydrates, saturated fat and sodium has increased in the daily diet. These types of foods are poor in vitamins, minerals, complex carbohydrates and fibres<sup>(6–8)</sup> and contribute to the increase in the prevalence of overweight and other cardiometabolic risk factors also among children<sup>(8,9)</sup>.

Thus, it is important to identify inappropriate eating habits in childhood to intervene early and prevent the occurrence of lifelong illness.

Recent studies have evaluated food consumption in the child population using the analysis of dietary patterns<sup>(7,10)</sup>, because individuals do not consume nutrients separately, but rather meals composed of a wide variety of nutrients, which undergo interactions. In this way, some studies have pointed out that socio-economic characteristics such as income and maternal education are predictors of a child’s food consumption, but there is no consensus on this relationship between studies<sup>(11–14)</sup>. In addition, there is evidence that breast-feeding duration may influence the development of eating habits in childhood<sup>(15)</sup>. However, most of these studies evaluated the effect of breast-feeding on the

\*Corresponding author. Email sarahvieiraufv@gmail.com

consumption of specific food groups rather than on overall food consumption<sup>(2,5)</sup>.

In this context, the present study aimed to identify the dietary patterns of children aged 4–7 years and verify their association with sociodemographic characteristics, lifestyle habits and duration of exclusive breast-feeding (EBF).

## Materials and methods

The present study was retrospective analysis conducted in a cohort of children born in the maternity hospital in Viçosa, Minas Gerais, Brazil. Children were followed up by the Lactation Support Program (PROLAC) during their first year of life. PROLAC is a Program of the Federal University of Viçosa in partnership with the São Sebastião Hospital and the Human Milk Bank of the municipality. The programme's main activities are to provide guidelines for mothers in the postpartum period, aiming to promote breast-feeding and nutritional care to nursing mothers and children in the first year of life.

We selected PROLAC care records for the recruitment of children based on two inclusion criteria: (i) those containing information that allowed location of the child; and (ii) child's date of birth compatible with age between 4 and 7 years at the time of study. Considering these criteria, 669 children were eligible to participate. After at least three attempts to contact children through home visits, 176 were not located (change of address), seventy-five parents did not authorize participation or did not complete all stages of the study, and eight had health problems that prevented them from participating. In addition, seven children were excluded for incomplete food consumption data. Thus, 266 losses were recorded (39.8%) and the final sample comprised 403 children.

The parent/guardian answered a semi-structured questionnaire to provide sociodemographic information on gender, age, area of residence (urban/rural), maternal education, per capita income, work outside the home (yes/no) and maternal civil status (with/without partner). Maternal education and income were categorized into tertiles for the analysis. Information on EBF ( $\geq 4$  months and  $< 4$  months) was collected from PROLAC care records.

Information on lifestyle habits was obtained by applying a questionnaire adapted from Andaki<sup>(16)</sup>. The variables evaluated were daily screen time (television, computer, games console) categorized according to the American Academy of Pediatrics' recommendation<sup>(17)</sup> ( $< 2$  h and  $\geq 2$  h) and regular physical activity (yes/no).

We obtained information regarding food consumption by evaluating three food records filled out by the parent/guardian of the child during non-consecutive days, with one day at the weekend. All records were checked and reviewed by the researchers and the parent/guardian to reduce errors when filling out the record form. The data

were entered and processed using DietPro<sup>®</sup> software version 5i.

To assess the dietary patterns, the food items reported in the records were entered with their respective amounts in grams and each child's average consumption of each food item during the three days of investigation was calculated. Afterwards, the foods and preparations were grouped based on their nutritional characteristics or botanical composition<sup>(14,18,19)</sup>, resulting in nineteen food groups (Table 1). Foods or preparations consumed by less than 10% of the children were redistributed into groups with similar nutritional content.

## Data analysis

The identification of dietary patterns was performed *a posteriori* by principal component analysis (PCA), which allows food groups to be combined based on the correlations between them. Before proceeding to PCA, we evaluated the adequacy of the sample size, considering that the ratio of cases to food groups should be  $\geq 5$ <sup>(20)</sup>. The ratio found in the present study was 21.2 (403 children/19 food groups).

To assess the applicability of PCA, the Kaiser–Mayer–Olkin coefficient was estimated and Bartlett's sphericity test was performed; the Kaiser–Mayer–Olkin coefficient for each variable should be  $\geq 0.5$ <sup>(20)</sup> and Bartlett's test of sphericity should be significant. We found a Kaiser–Mayer–Olkin coefficient of 0.561 and  $P < 0.001$  for Bartlett's test. We also assessed the adequacy of each variable (food group) for PCA, evaluating the anti-image correlation matrix.

To improve interpretation of the factors, we applied orthogonal varimax rotation, which allows the resulting factors not to be correlated<sup>(21)</sup>. The number of factors to be retained was determined based on the Catell (scree plot) chart, where the points of greatest slope indicate the number of factors to be considered in the analysis<sup>(21)</sup>. The interpretability of factor loads was also considered to select the number of components.

Foods or food groups with an absolute factor loading of  $\geq 0.25$  were considered to be strongly associated with the component, providing better information for the identification of a dietary pattern. Food groups with a positive factor loading contribute directly to a dietary pattern, while those with a negative factor loading are inversely associated with the pattern<sup>(21)</sup>. When a food was present in more than one pattern, it was maintained in the pattern where it presented the highest factor loading.

A food/food group that presented an absolute factor loading of  $\geq 0.25$  in more than one component was standardized to be kept in the group with the highest factor loading. For the designation of dietary patterns, we considered the characteristics of the foods/food groups that contributed the most to each pattern, as well as the nomenclatures used in the literature<sup>(14,22,23)</sup>.

**Table 1** Foods and food groups, categorized according to nutritional characteristics or botanical composition, consumed by children aged 4–7 years (*n* 403), Viçosa, Minas Gerais, Brazil, 2015–2016

Food/food group	Foods reported in the food records
Milk and derivates	Whole milk, cheeses, curd cheese, yoghurt, petit suisse cheese, fermented milk, dairy drinks
Chocolate and sugar	Cocoa powder and sugar
Coffee and mate tea	Coffee and mate tea
Butter and margarine	Butter and margarine
Bread, cakes and cookies	French bread, toast bread, toast, maize biscuit, water and salt biscuit, doughnut, biscuit, potato biscuit, buttery biscuit, cereal, plain cake, corn cake, chocolate cake
Filled cookies	Filled cookies
Beans	Cooked beans, beans <i>tutu</i> , <i>tropeiro</i> beans, <i>feijoada</i> stew
White rice	Cooked white rice
Leafy vegetables	Lettuce, cabbage, mustard, endive, chard, spinach, Barbados gooseberry, broccoli, cauliflower, parsley, chives, watercress, arugula, corm from elephant-eye plant, cabbage
Vegetables	Tomato, onion, cucumber, peppers, zucchini, pumpkin, eggplant, okra, scarlet eggplant, carrot, beet, pod, chayote, vegetable mayonnaise, vinaigrette
Tubers, polenta and flours	Baked potato, mashed potatoes, yams, arracacha, cooked cassava, sweet potatoes, cooked polenta, polenta, toasted cassava flour ( <i>farofa</i> ), cassava flour, corn flour
Fruits	Banana, apple, orange, clementine, pear, guava, papaya, soursop, avocado, plum, peach, Barbados cherry, mango, grapefruit, watermelon, melon, strawberry, pineapple, passion fruit, star fruit, lemon, siriguela, blackberry, jabuticaba, fruit salad
Natural juice	Orange, lemon, pineapple, watermelon, acerola, guava, passion fruit, mango, grape, strawberry, apple, papaya
Artificial juice and soft drinks	Juice powder, can juice, concentrated juice, soda, cola
Meat, fish and eggs	Cooked beef, fried beef, roasted pork, fried pork, fried fish, cooked fish, cod, sardines, chicken heart, chicken gizzard, beef liver and chicken liver, boiled egg, fried egg, omelette
Fried foods, snacks and sausages	Fried potato, fried cassava, chicken <i>cozinha</i> , kibbeh, hot dog, fried <i>pastel</i> , pot pie, cheese bread, pasty, chicken pie, leafy pastry, fried nugget, hamburger, pizza, salted popcorn, potato chips, ham, mortadella, sausage, bacon
Sweets	Chocolate, ice cream, popsicle, condensed milk, brigadeiro, party candy, dulce de leche, peanut candy, cake topped with filling, sweet popcorn, sweet rice, sweet canjica, candies, chewing gum, gelatine
Broths and soups	Vegetable soup, vegetable soup and pasta, <i>canjiquinha</i> , cabbage porridge, chicken soup, green broth, cassava broth
Pasta	Gnocchi, lasagne, meat pancakes, boiled pasta, garlic and oil pasta, pasta noodles, bolognaise pasta, instant noodles

After the identification of dietary patterns, we calculated the factor scores for each child in the study. Thus, each child had a factor score in all identified patterns, with a position based on the degree of participation in each pattern<sup>(18,22)</sup>.

Statistical analyses were performed using the statistical software package Stata version 13.0, with normality of the variables evaluated by the Shapiro–Wilk test. The children’s consumption scores for each dietary pattern were categorized into tertiles for the association analysis. Simple multinomial logistic regression was applied to evaluate the association between the predictive variables (sociodemographic characteristics, lifestyle habits and EBF) and each dietary pattern (dependent variables). The variables with  $P < 0.20$  in the bivariate analysis were included in the multivariate logistic regression models to estimate OR and 95% CI. The final model considered the variables that were associated with the dependent variables at  $P < 0.05$ .

The study was approved by the Ethics Committee for Research with Human Beings of the Federal University of Viçosa (reference number 892476/2014). The children’s participation occurred after the parents or guardians signed the Term of Free and Informed Consent.

## Results

The distribution of the sociodemographic, lifestyle and EBF variables in the sample is described in Table 2. More than half of the children evaluated were male (55%) with age between 6 and 7 years (55%). We found that the majority of the children (61%) received EBF until at least 4 months of life, and most (71%) children reported screen time of  $>2$  h/d. Regarding maternal education, 51% of mothers reported 8–11 years of schooling.

Table 3 shows the foods/food groups and energy consumed by the studied children. High median values were found for the consumption of milk and milk products (281 g/d) and artificial juice and soft drinks (110 g/d). However, lower consumption medians were found for leafy vegetables (7 g/d), vegetables (18 g/d) and natural juices (28 g/d).

PCA extracted five dietary patterns, which explained 42% of the variance in the data (Table 4). The first dietary pattern, named ‘Traditional’, explained 10% of the variance and consisted of typical foods/food groups present in the Brazilian diet, including: white rice; beans; vegetables; tubers, polenta and flours; and meat, fish and eggs. The ‘Unhealthy’ pattern was represented mainly by

**Table 2** Distribution of sociodemographic characteristics, lifestyle habits and breast-feeding in the sample of children aged 4–7 years (*n* 403), Viçosa, Minas Gerais, Brazil, 2015–2016

Variable	<i>n</i>	%
Gender		
Male	222	55.1
Female	181	44.9
Age		
4–5 years	179	44.4
6–7 years	224	55.6
Maternal work outside the home*		
Yes	283	70.6
No	118	29.4
Maternal education†,‡		
< 8 years	136	39.1
8–11 years	202	50.6
> 11 years	61	15.3
Monthly per capita income†,§		
R\$ 67.66–224.99	133	33.4
R\$ 225.00–340.00	154	38.7
R\$ 340.01–2500.00	111	27.9
Maternal civil status*		
With partner	321	80.1
Without partner	80	19.9
Residential area		
Urban	370	91.8
Rural	33	8.2
EBF until 4 months		
Yes	241	60.9
No	135	39.1
Screen time¶		
≤ 2 h/d	116	29.0
> 2 h/d	284	71.0
Regular practise of physical activity		
Yes	64	15.9
No	339	84.1

EBF, exclusive breast-feeding (defined as baby receives only breast milk, straight from the mother, or human milk from another source, without any other product or solids).

\**n* 401.

†Categorization in tertiles.

‡*n* 399.

§*n* 398.

||*n* 396.

¶*n* 400.

foods/food groups with high sugar and fat content, such as: artificial juice and soft drinks; fried foods, snacks and sausages; sweets; and stuffed cookies. In addition, the food group broths and soups was negatively (inversely) associated in this pattern.

The 'Milk and chocolate' pattern was so named because it was mainly represented by milk and milk products, and chocolate and sugar, foods that were highly frequently consumed by the children studied. The group coffee and tea was inversely related to this food pattern, indicating that children who consumed milk with chocolate were not in the habit of drinking coffee and tea. The 'Snack' dietary pattern was represented mainly by typical bakery foods, such as: breads, cakes and cookies; butter and margarine; and coffee and tea. The foods/food groups stuffed cookies, fruits and pasta were negatively associated with this pattern. The last pattern, entitled 'Healthy', was represented mainly by natural juice, fruits, vegetables, and broths and soups.

In the bivariate logistic regression analysis (data not shown), children with higher per capita income and

**Table 3** Consumption of foods/food groups and energy by children aged 4–7 years (*n* 403), Viçosa, Minas Gerais, Brazil, 2015–2016

Food/food group (g/d)	Median	Q1–Q3
Milk and derivatives	280.8	187.5–428.3
Artificial juice and soft drinks	110.0	0–220.0
Beans	99.6	65.0–139.3
White rice	100.0	75.0–136.6
Meat, fish and eggs	63.3	43.3–88.6
Bread, cakes and cookies	58.3	36.3–83.6
Fruits	60.0	10.0–126.7
Fried foods, snacks and sausages	40.0	18.3–75.0
Natural juice	27.5	0–110.0
Coffee and tea	21.6	0–57.5
Vegetables	17.5	5.3–35.8
Tubers, polenta and flours	20.0	5.0–39.1
Sweets	13.6	0–43.3
Pasta	13.3	0–36.6
Leafy vegetables	6.6	0–14.0
Chocolate and sugar	10.0	0–61.3
Butter and margarine	1.3	0–3.3
Filled cookies	0.0	0–16.6
Broths and soups	0.0	0–30.0
Energy (kJ/d)	6252.2	5382.3–7265.5
Energy (kcal)	1494.3	1286.2–1738.5

Q1, quartile 1 (25th percentile); Q3, quartile 3 (75th percentile).

mothers with higher education showed greater adherence to the 'Unhealthy', 'Milk and chocolate' and 'Healthy' patterns ( $P < 0.005$ ). Children with screen time of  $> 2$  h/d ( $P = 0.003$ ) and those aged 6 and 7 years ( $P = 0.001$ ) were more likely to consume more 'Unhealthy' foods. The older children (aged 6 and 7 years) showed lower adherence to the 'Milk and chocolate' pattern, compared with children aged 4 and 5 years ( $P = 0.024$ ). Children who received EBF until at least 4 months of life had moderate adherence (second tertile) to the 'Snack' pattern ( $P = 0.026$ ).

Table 5 shows the final models of the multivariate logistic regression analysis for each dietary pattern (dependent variables) and sociodemographic characteristics, lifestyle habits and EBF (independent variables). As in the simple model, no variable was associated with the 'Traditional' pattern. Even after adjustment, children with higher per capita income showed higher adherence to the 'Unhealthy', 'Milk and chocolate' and 'Healthy' patterns ( $P < 0.05$ ).

Older children (aged 6 and 7 years) had a 2.25 times greater chance of consuming 'Unhealthy' foods than children aged 4 and 5 years. In addition, children who did not receive EBF until at least 4 months of life also had higher adherence to this pattern (OR = 1.71; 95% CI 1.02, 2.86). After adjustment in the multivariate model, the association between lower EBF duration and moderate food consumption of the 'Snack' pattern was maintained (OR = 1.76; 95% CI 1.06, 2.90).

## Discussion

The present study identified five dietary patterns ('Traditional', 'Unhealthy', 'Milk and chocolate', 'Snack' and

**Table 4** Dietary patterns and factor loadings of foods/food groups consumed by children aged 4–7 years (*n* 403), Viçosa, Minas Gerais, Brazil, 2015–2016

Food/food group	Dietary pattern*				
	Traditional	Unhealthy	Milk and chocolate	Snack	Healthy
Milk and derivates	−0.038	−0.234	<b>0.738</b>	0.164	−0.018
Chocolate and sugar	−0.026	0.028	<b>0.856</b>	0.028	−0.050
Coffee and tea	0.036	−0.234	<b>−0.502</b>	<b>0.252</b>	−0.181
Butter and margarine	0.194	0.182	0.034	<b>0.573</b>	−0.027
Bread, cakes and cookies	0.044	0.032	−0.012	<b>0.703</b>	0.091
Stuffed cookies	−0.111	<b>0.273</b>	0.105	<b>−0.381</b>	−0.026
Beans	<b>0.684</b>	−0.159	−0.102	−0.139	0.110
White rice	<b>0.749</b>	0.004	−0.144	−0.043	−0.161
Leafy vegetables	<b>0.499</b>	−0.051	0.073	0.178	0.256
Vegetables	0.371	0.031	0.002	0.055	<b>0.433</b>
Tubers, polenta and flours	<b>0.381</b>	−0.080	0.017	0.204	−0.019
Fruits	0.125	−0.114	0.115	<b>−0.273</b>	<b>0.532</b>
Natural juice	−0.132	0.034	−0.051	0.231	<b>0.620</b>
Artificial juice and soft drinks	−0.127	<b>0.755</b>	0.020	0.001	−0.083
Meat, fish and eggs	<b>0.408</b>	0.329	0.093	0.122	−0.142
Fried foods, snacks and sausages	−0.088	<b>0.631</b>	−0.106	−0.028	0.067
Sweet	0.010	<b>0.477</b>	0.031	−0.144	0.443
Broth and soups	−0.090	<b>−0.334</b>	−0.021	0.049	<b>0.306</b>
Pasta	0.052	0.151	−0.072	<b>−0.458</b>	0.027
% of variance explained	9.8	9.0	8.5	8.0	7.0
Total variance explained (%)	42.3				

\*Dietary patterns were extracted by principal component analysis with varimax rotation and Kaiser normalization, and named through main components. Bold values indicate absolute factor loadings  $\geq 0.25$ .

**Table 5** Final models of the multinomial logistic regression analysis for the association of socio-demographic characteristics, lifestyle habits and breast-feeding with tertiles of consumption of dietary patterns (outcomes) among children aged 4–7 years (*n* 403), Viçosa, Minas Gerais, Brazil, 2015–2016

	OR	95% CI	P value
Unhealthy pattern*			
Children aged 6 and 7 years	2.25	1.34, 3.77	0.002
Third tertile of per capita income	2.47	1.35, 4.54	0.003
EBF for <4 months	1.71	1.02, 2.86	0.040
Milk and chocolate pattern†			
Third tertile of per capita income	2.84	1.48, 5.45	0.002
Snack pattern‡ (second tertile of consumption scores)			
EBF for <4 months	1.76	1.06, 2.90	0.027
Healthy pattern§			
Third tertile of per capita income	2.94	1.43, 6.05	0.003

EBF, exclusive breast-feeding.

\*Adjusted for gender, age, per capita income, maternal work, screen time and EBF duration.

†Adjusted for age, per capita income, maternal work, civil status and physical activity practise.

‡Adjusted by for age, maternal education, civil status, physical activity practise and EBF duration.

§Adjusted for gender, age, per capita income and physical activity practise.

‘Healthy’) that represented the overall dietary habits of the children evaluated and explained 42% of the original data variance. Other studies carried out to determine the dietary patterns in Brazilian children showed similarities with our study. Most of them identified a dietary pattern composed mainly of foods that are part of the population’s eating habits such as rice, beans, tubers and meat, commonly called ‘Traditional’. The other characteristic patterns found in the studies showed the consumption of foods rich in refined sugars, fats and sodium, named the ‘Unhealthy’ pattern in our study, and a ‘Healthy’ pattern which is based on foods such as fruits, vegetables and natural juices<sup>(14,23,24)</sup>.

In another study carried out in Viçosa, Minas Gerais, Brazil, Villa *et al.*<sup>(14)</sup> identified among children aged 8 and 9 years a dietary pattern composed of milk and chocolate, which they named as ‘Monotone’, confirming that the consumption of milk added by chocolate is part of the eating habits of the children in the municipality. These authors found that the mean consumption of chocolate was 18.7 g/d, above the mean found in our study (14 g/d). Another study carried out in southern Brazil with children aged 1–6 years also identified a pattern containing milk and chocolate. The authors suggested that children who adhere to this pattern probably do not consume main meals properly, replacing them with quick snacks, such as chocolate milk<sup>(24)</sup>.



The 'Snack' pattern identified in the present study was also observed in pre-school children from Diamantina, Minas Gerais, Brazil, based mainly on bakery products and butter/margarine<sup>(23)</sup>. A cohort study of Chinese children aged 13 months and 6 years identified a food pattern composed mainly of refined carbohydrates and confectionery products<sup>(25)</sup>.

In fact, the food patterns identified in the present study, except the 'Healthy' one, were in line with the worldwide trend of frequent consumption of ultra-processed foods by children that are high in fat and refined carbohydrates, such as baked goods, sausages, sweetened beverages and sweets<sup>(14,23,26)</sup>. The excessive consumption of these foods in association with a sedentary lifestyle is related to risk of developing overweight and cardiometabolic diseases in the child population<sup>(5,25,27,28)</sup>.

An interesting result of our study was the association of income with the dietary patterns 'Unhealthy' and 'Milk and chocolate', composed mainly of foods high in fat and refined sugars. A similar result was observed in Nobre *et al.*'s study<sup>(23)</sup> in Brazil, in which children belonging to higher-income families had higher participation in the pattern based on unhealthy foods, such as sweets, sodas and stuffed cookies. However, in a study carried out in Spain, low-income children and adolescents were more likely to consume unhealthy foods<sup>(29)</sup>.

Higher consumption of the 'Healthy' pattern was also associated with higher per capita income. This result corroborates the results of a multicentre study with children aged 2–9 years in Europe, where children whose parents had higher income and schooling consumed more healthy foods<sup>(12)</sup>. However, the authors emphasize that the average consumption of foods belonging to this pattern, such as fruits and vegetables, was low. We also observed low average consumption of these foods among the children assessed, mainly leafy vegetables and natural juices. While the mean consumption of artificial juices and soft drinks was 148 ml/d, the consumption of natural juices was 71 ml/d, evidencing the trend of consumption of industrialized sweetened drinks by children which has been associated with unfavourable health outcomes<sup>(4,8)</sup>.

No sociodemographic, lifestyle or breast-feeding variables were related to the 'Traditional' dietary pattern after adjustment of the multivariate model. Thus, it is suggested that regardless of income, EBF duration and lifestyle, most of the children frequently consumed the 'Traditional' dietary pattern, which is composed of the typical Brazilian foods such as rice and beans. However, other studies have found an association between food consumption of the 'Traditional' dietary pattern and household income. Villa *et al.*<sup>(14)</sup> observed that children of lower socio-economic level showed greater adherence to this pattern. However, other works identified an inverse association of the 'Traditional' pattern with lower socio-economic level<sup>(12,24)</sup>.

Although maternal work outside the home was not associated with any of the dietary patterns in our study,

most children with the highest per capita income were those whose mothers worked. Thus, it is believed that the inclusion of women in the labour market makes it difficult to prepare meals, which leads to greater consumption of foods outside the home, especially industrialized foods<sup>(27)</sup>.

The bivariate analysis showed that children with screen time of more than 2 h/d presented higher intake of foods from the 'Unhealthy' pattern, but after adjusting for other variables in the multivariate model, the association was not maintained. In a study conducted with American schoolchildren, it was found that those with greater daily time in front of a television and games console were more likely to consume fast foods and sugary drinks<sup>(13)</sup>. The broadcasting and encouragement of television advertising for processed food products, especially ultra-processed products, is a factor that influences the high consumption of these foods by children, especially those who spend most of their free time in sedentary activities<sup>(30)</sup>.

The current study presents novel results on the relationship of EBF for less than 4 months with the consumption of foods of the 'Unhealthy' and 'Snack' patterns, after adjustment, in the multivariate models. Apparently, only one study has previously used PCA to investigate the influence of EBF during the early months of life on childhood food patterns<sup>(31)</sup>, showing that EBF duration was positively associated with the 'Healthy' pattern. These authors found associations between breast-feeding only and the 'Healthy' pattern, to which breast-fed children showed greater adherence. Perrine *et al.*<sup>(2)</sup>, who evaluated the association of EBF with the consumption of certain food groups by 6-year-olds, identified that breast-feeding duration was inversely associated with consumption of sugary drinks. Soldateli *et al.*<sup>(3)</sup> found no association between lactation time and consumption of fruits and vegetables by children at 4–7 years of age.

Infant dietary behaviour develops early and is influenced by different factors, including maternal diet during the intra-uterine phase<sup>(32)</sup>. Breast-fed children can taste different flavours of the foods consumed by the mother. This provides greater variation of the palate, when compared with children fed formula. Thus, studies have shown that children who receive breast milk have better acceptance of new foods, especially fruits and vegetables, and dietary habits tend to remain in later stages of life<sup>(1,32)</sup>.

As a possible limitation in the present study, we emphasize the subjectivity during the identification of food patterns in several stages of the analysis, such as the criteria used to group foods, the number of factors to be retained and in the way the identified patterns are named. However, to minimize the effects of this limitation, the criteria adopted during all stages of the analysis were described in detail. As a positive point, we highlight the use of three food records for evaluation of consumption, which reduces the intra-individual variability of intake and allows us to investigate the habitual food consumption of the children. In addition, the use of recorded data





on breast-feeding duration is emphasized, avoiding memory bias.

## Conclusion

In conclusion, we identified five dietary patterns in children aged 4–7 years. The highest per capita income was associated with high consumption of foods from the 'Unhealthy', 'Milk and chocolate' and 'Healthy' patterns. Children who were not exclusively breast-fed until 4 months of life showed more adherence to the 'Unhealthy' and 'Snack' patterns, and older children aged 6 and 7 years also consumed more foods of the 'Unhealthy' pattern.

In view of the identified food patterns, most of them based mainly on foods high in sugars, fats and sodium, it is necessary to implement public food and nutrition policies directed to children, since eating habits are established during childhood and when inadequate and maintained throughout life, they can lead to the development of diseases. In addition, the importance of support and encouragement of EBF during the first months of life is emphasized, which, in addition to the many benefits already confirmed, can positively influence lifelong eating habits.

## Acknowledgements

**Acknowledgements:** The authors thank the National Council for Scientific and Technological Development (CNPq) and the Foundation for Research Support of the State of Minas Gerais (FAPEMIG) for the financial support; the Coordination for the Improvement of Higher Education Personnel (CAPES) for granting a scholarship; and the children, parents and guardians for participating in the study. **Financial support:** This work was supported by CNPq (grant number 485124/2011-4) and FAPEMIG (grant number 02055-13). CNPq, FAPEMIG and CAPES had no role in the design, analysis or writing of this article. **Conflict of interest:** The authors declare that they have no conflicts of interest. **Authorship:** S.A.V. assisted the conception and design of this work, assisted the data collection, analysis and interpretation of the data, conducted the literature search, as well as wrote the manuscript. P.C.A.F. and C.S.A. assisted the conception and design of this work, assisted the data collection, analysis and interpretation, as well as revised and approved the final version to be published. H.H.M.H., A.Q.R., P.F.P. and S.E.P. revised and approved the final version to be published. S.C.C.F. designed the study including the data collection, coordinated and supervised and approved the final version to be published. **Ethics of human subject participation:** This study was conducted according to the guidelines laid down in the Declaration of Helsinki and

all procedures involving human subjects were approved by the Ethics Committee on Human Research of the Federal University of Viçosa (case number 892476/2014). All participants, as well as their responsible adults, were informed about the objectives of the research and informed consent was obtained from all children's parents/guardians.

## References

1. Vitolo MR, Rauber F, Campagnolo PD *et al.* (2010) Maternal dietary counseling in the first year of life is associated with a higher healthy eating index in childhood. *J Nutr* **140**, 2002–2007.
2. Perrine CG, Galuska DA, Thompson MPH *et al.* (2014) Breastfeeding duration is associated with child diet at 6 years. *Pediatrics* **134**, 50–55.
3. Soldateli B, Vigo A & Giugliane ERJ (2016) Effect of pattern and duration of breastfeeding on the consumption of fruits and vegetables among preschool children. *PLoS One* **11**, e0148357.
4. Dishchekenian VRM, Escrivão MAMS, Palma D *et al.* (2011). Dietary patterns of obese adolescents and different metabolic effects. *Rev Nutr* **24**, 17–29.
5. Zarrinpar A, Chaix A & Panda S (2016) Daily eating patterns and their impact on health and disease. *Trends Endocrinol Metab* **27**, 69–83.
6. Brazilian Institute of Geography and Statistics (2010) *Survey of Family Budgets 2008–2009. Anthropometry and Nutritional Status of Children, Adolescents and Adults in Brazil*. Rio de Janeiro, RJ: IGBE.
7. Kiefe de Jong JC, Vries JH *et al.* (2013) Socio-demographic and lifestyle determinants of 'Western-like' and 'Health conscious' dietary patterns in toddlers. *Br J Nutr* **109**, 137–147.
8. Rinaldi AEM, Gabriel GFPC, Moreto F *et al.* (2016) Dietary factors associated with metabolic syndrome and its components in overweight and obese Brazilian schoolchildren: a cross-sectional study. *Diabetol Metab Syndr* **8**, 58.
9. Sparrenberger K, Friedrich RR, Schiffner MD *et al.* (2015) Ultra-processed food consumption in children from a Basic Health Unit. *J Pediatr (Rio J)* **91**, 535–542.
10. Voortman T, Leermakers ETM, Franco OH *et al.* (2016) *A priori* and *a posteriori* dietary patterns at the age of 1 year and body composition at the age of 6 years: the Generation R Study. *Eur J Epidemiol* **31**, 775–783.
11. Silva RC, Assis A.M, Szarfarc SC *et al.* (2012) Socioeconomic inequality shaping the dietary patterns of children and teens. *Rev Nutr* **25**, 451–461.
12. Fernandez-Alvira JM, Rnhorst CB, Bammann K *et al.* (2015) Prospective associations between socio-economic status and dietary patterns in European children: the Identification and Prevention of Dietary- and Lifestyle-induced Health Effects in Children and Infants (IDEFICS) Study. *Br J Nutr* **113**, 517–525.
13. Lowry R, Michael S, Demissie Z *et al.* (2015) Associations of physical activity and sedentary behaviors with dietary behaviors among US high school students. *J Obes* **2015**, 876524.
14. Villa JKD, Silva AR, Santos TSS *et al.* (2015) Dietary patterns of children and socioeconomical, behavioral and maternal determinants. *Rev Paul Pediatr* **33**, 302–309.
15. Maier AS, Chabanet C, Schaal B *et al.* (2008) Breastfeeding and experience with variety early in weaning increase infants' acceptance of new foods for up to two months. *Clin Nutr* **27**, 849–857.



16. Andaki ACR (2010) Anthropometry and level of physical activity in predicting metabolic changes in 10-year-old children. Master's Thesis, Federal University of Viçosa.
17. American Academy Pediatrics, Committee on Public Education (2001) Children, adolescents, and television. *Pediatrics* **107**, 423–426.
18. Hu FB (2002) Dietary pattern analysis: a new direction in nutritional epidemiology. *Curr Opin Lipidol* **13**, 3–9.
19. Matos SMA, Barreto ML, Rodrigues LC *et al.* (2014) Dietary patterns of children under five years of age living in the State capital and other counties of Bahia State, Brazil, 1996 and 1999–2000. *Cad Saude Publica* **30**, 44–54.
20. Hair JF, Anderson RE, Tatham RL *et al.* (2005) *Análise Multivariada de Dados*, 9th ed. Porto Alegre, RS: Bookman5.
21. Olinto MT (2007) Dietary patterns: principal components analysis. In *Epidemiologia Nutricional*, pp. 213–226 [G Kag, R Sichieri and DP Gigante, organizadores]. Rio de Janeiro, RJ: Fiocruz/Atheneu.
22. Salvatti AG, Escrivão MAMS, Taddei JAAC *et al.* (2011) Eating patterns of eutrophic and overweight adolescents in the city of São Paulo, Brazil. *Rev Nutr* **24**, 703–713.
23. Nobre L, Lamounier JA & Franceschini SCC (2012) Pre-school children dietary patterns and associated factors. *J Pediatr (Rio J)* **88**, 129–136.
24. Souza RLV, Madruga SW, Gigante DP *et al.* (2013) Dietary patterns and associated factors among children one to six years of age in a city in southern Brazil. *Cad Saude Publica* **29**, 2416–2426.
25. Hooven EHV, Heppel DHM, Kiefe de Jong JC *et al.* (2015) Infant dietary patterns and bone mass in childhood: the Generation R Study. *Osteoporos Int* **26**, 1595–1604.
26. World Health Organization (2003) *Diet, Nutrition and the Prevention of Chronic Diseases. Joint WHO/FAO Expert Consultation. WHO Technical Report Series* no. 916. Geneva: WHO.
27. Rinaldi AE, Pereira AF, Macedo CS *et al.* (2008) Feeding practices and physical inactivity contributions to childhood overweight. *Rev Paul Pediatr* **26**, 271–277.
28. Santos NHA, Fiaccone R, Barreto ML *et al.* (2014) Association between eating patterns and body mass index in a sample of children and adolescents in Northeastern Brazil. *Cad. Saude Publica* **30**, 2235–2245.
29. Aranceta J, Perez-Rodrigo C, Ribas L *et al.* (2003) Socio-demographic and lifestyle determinants of food patterns in Spanish children and adolescents: the enKid study. *Eur J Clin Nutr* **57**, Suppl. 1, 40–44.
30. Story M & Faulkner P (1990) The prime time diet: a content analysis of eating behavior and food messages in television program content and commercials. *Am J Public Health* **80**, 738–740.
31. Grieger JA, Scott J & Cobiac L (2011) Dietary patterns and breast-feeding in Australian children. *Public Health Nutr* **14**, 1939–1947.
32. Beauchamp GK & Mennella JA (2009) Early flavor learning and its impact on later feeding behavior. *J Pediatr Gastroenterol Nutr* **48**, Suppl. 1, 25–30.